

Recombinant Human IL-18

 Catalog#:AC13069 Derived from *E.coli*

DESCRIPTION	<p>Recombinant Human Interleukin-18 is produced by our E.coli expression system and the target gene encoding Tyr37-Asp193 is expressed.</p> <p>Accession#: Q14116</p> <p>Known as: Interleukin-18; Iboctadekin; Interferon gamma-inducing factor; IFN-gamma-inducing factor; Interleukin-1 gamma; IL-1 gamma; GIF; IL-18; IL-1g; IL1F4; MGC12320</p>
FORMULATION	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7 °C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20 °C for 3 months.</p>
RECONSTITUTION	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100 μg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass: 18.2 kDa AP Mol Mass: 17 kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
	<p>Interleukin-18 is a secreted protein and it belongs to the IL-1 family. IL-18 is a proinflammatory cytokine and produced by macrophages and other cells. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-γ (IFN-γ) or type II interferon that plays an important role in activating the macrophages or other cells.</p>
	<p>SDS-PAGE</p> 